



www.vompti.com

THORACIC SPINE CASE #1


Aaron Hartstein, PT, DPT, OCS, FAAOMPT
AJ Lievre, PT, DPT, OCS, CMPT

Orthopaedic Manual Physical Therapy Series
Charlottesville 2017-2018



Orthopaedic Manual Physical Therapy Series 2017-2018

Body Chart



Body Chart – Initial Hypothesis:

- T8/9 Segmental Dysfunction (Somatic)
- Intercostal Sprain/Strain
- T8/9 Radiculopathy
- Visceral Referral/Red Flag?
 - Gallbladder?
 - Liver?




Orthopaedic Manual Physical Therapy Series 2017-2018


www.vompti.com

Subjective History

- 43 y/o R handed male with thoracic spine and ribcage symptoms
- 2 wk history of thoracic symptoms after shoveling and moving 4 tons of gravel which took 5 hours
- Mid-back “ache” described towards end of shoveling and with difficulty sleeping that night
- Worsening in last 2 weeks with increased irritability
- 1st episode of thoracic area symptoms other than gallbladder “attack”
- Reports hx of low level but dull R lower quadrant symptoms, at times after he eats certain foods
- PMH significant for gallbladder dysfunction, gallstones and previous alcohol abuse



Subjective *Asterisks* Signs/Symptoms: (Aggravating/Easing factors, Description/location of symptoms, Behavior, Mechanism of injury):



- Symptom Behavior:
 - Constant, deep R thoracic ache/burning which intermittently radiates P-A
- Symptoms can occur together but appear unrelated
- Can have posterior thoracic pain without radiation anterior-laterally
- Currently still working as stone mason
- Aggs: Deep breath, twisting, cough/sneeze, reaching down towards floor, rolling in bed at night, certain foods
- Eases: changing position, rest, pillow under R arm, L SL with arm overhead
- Somewhat activity/positional dependent and worsens throughout day



STRUCTURE at Fault:				
Joints in/refer to the painful region	Myofascial tissue in/refer to the painful region	Non Contractile tissue in/refer to the painful region	Neural tissue in/refer to the painful region	Other structures that must be examined – non MSK
C7-T2 and T8/9 Z, CV, CT Jts T8/9 Costochondral Jt	T8/9 paraspinals and multifidus, intercostals, lower trap	T8/9 Capsule/IVD Ribs (Fx)	T8/9 nn root	Visceral: Gallbladder Liver Lung Ankylosing-Spondylitis

Primary HYPOTHESIS after Subjective Examination: – **T8/9 mechanical dysfunction (CV/CT) with somatic referral**

Differential List (Rank/List in order to rule out):
Visceral Referral, Cervical Somatic Dysfunction, Muscular Strain

Orthopaedic Manual Physical

SPINE Volume 18, Number 7, pp 807-811
©1994, J. B. Lippincott Company

Thoracic Zygapophysial Joint Pain Patterns

A Study in Normal Volunteers



Paul Dreyfuss, MD,*† Claire Tibiletti, MD,* and Susan J. Dreyer, MD‡§

Figure 3. A composite map of the results in all volunteers showing referral patterns from the T3-T4 to T10-T11 thoracic zygapophysial joints.

Orthopaedic Manual Physical Therapy series 2017

Thoracic Zygapophyseal Joint Referral Patterns C7/1-T2/3 and T11/12

- Fukui et al.



Research article

Open Access

Thoracic costovertebral joint pain patterns: a study in normal volunteers

Brian A Young^{*1}, Howard E Gill², Robert S Wainner³ and Timothy W Flynn⁴




Figure 4 Composite diagram of costovertebral joint pain patterns.

Thoracic Pain Patterns

- Discogenic: complaint of central posterior pain which goes through the sternum. Often described as being “underneath sternum”
- Costovertebral, Costotransverse, or Z-Joint: complaint of a horizontal or lateral spread of pain
- Nerve Root: Pain around the line of the rib (T1 nerve root may give arm symptoms and pain across inferior angle of scapula) (Rule Out - Herpes Zoster Virus)
- Costochondral Joint: anterior chest pain (over the joint)
- Must rule out cervical referral to thoracic spine and scapular region (Facet/Disc)



Identify any potential risk factors (Yellow, Red flags, non MSK involvement, biopsychosocial)

- **History of Gallbladder dysfunction and alcohol abuse**
- **with potential non MSK/visceral referral**
- Must rule out serious pathological or visceral cause of symptoms
- Since presence of primary thoracic pain is low, 15%, must be suspicious of non-mechanical causes with thoracic spine and chest pain
- Visceral sources considered when no clear mechanical features exist
 - Myocardial ischemia, AAA, peptic ulcer, acute cholecystitis, renal colic, pyelonephritis
 - Majority of visceral organs innervated by T/S spinal nerves



Thoracic Spine Differential Diagnosis



OR



018

www.vompti.com

[RESIDENT'S CASE PROBLEM]

JASON R. RODEGHERO, PT, DPT, OCS, ATC, FAAOMPT^{1,2,3} • THOMAS R. DENNINGER, PT, DPT, OCS, FAAOMPT⁴ • MICHAEL D. ROSS, PT, DHS⁵

Abdominal Pain in Physical Therapy Practice: 3 Patient Cases

- 46y/o female R groin & R lower abdominal P!
- Insidious onset x 6 weeks, deep dull ache with cramping
- Aggs: none
- Eases: NSAIDs
- B & B (-), excessive menstrual bleeding last 2-3 months
- No to all 5 questions in cluster
- ROM, Neuro, Provocation testing (-)
- (+) R LQ palpation

TABLE 2

ABDOMINAL PAIN OF MUSCULOSKELETAL ORIGIN QUESTION CLUSTERS*

Cluster 1:

1. "Does coughing, sneezing, or taking a deep breath make your pain feel worse?" (yes)
2. "Do activities such as bending, sitting, lifting, twisting, or turning over in bed make your pain feel worse?" (yes)
3. "Has there been any change in your bowel habit since the start of your symptoms?" (no)

Cluster 2:

1. "Does eating certain foods make your pain feel worse?" (no)
2. "Has your weight changed since your symptoms started?" (no)

*Answering yes to either of the first 2 questions and no to the third question in cluster 1 results in a moderate probability that the patient's abdominal complaints are of musculoskeletal origin. The probability increases to strong if both questions in cluster 2 are answered with a no.


Sparkes V, Prevost AT, Hunter JO. Derivation and identification of questions that act as predictors of abdominal pain of musculoskeletal origin. *Eur J Gastroenterol Hepatol.* 2003;15:1021-1027. <http://dx.doi.org/10.1097/01.meg.0000059173.46867.Dc>




Orthopaedic Manual Physical Therapy Series 2017-2018


www.vompti.com

Visceral Referral Pattern



Abdominal Aortic Aneurysm





Medical Screening

- **Other Red Flags in T/S – infection, fracture, neoplasms and inflammatory disorders**
 - Spinal metastases (usually breast, lung, or colon primary) are most common forms of cancer in thoracic spine (Primary tumors rare)
 - Ankylosing Spondylitis – affects thoracic spine and rib joints with limited ribcage and chest expansion (hallmark is less than 2.5 cm)
 - AM Stiffness, sacroilitis, peripheral joint involvement, M>F 3:1, 15-40 y/o
 - Fractures – traumatic or osteoporotic
 - Men OR Women 60 or older presenting with acute thoracic spine pain must rule out




Visceral Palpation and Assessment


Thoracic-Specific Red Flags

- **Category I Findings**

- Viscerosomatic Pain
- Tumors and Fractures



http://www.wehelpwhat hurts.homestead.com/visceral-somatic_referral_patterns_resize_smaller.jpg



om

Murphy's Sign for Cholecystitis

- Sensitivity = 97
- Specificity = 48
- (+) LR = 1.9
- (-) LR = .06



(-)



Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Liver Palpation



(-)

Study	Reliability	Sensitivity	Specificity	LR+	LR-	QUADAS Score (0-14)
Halpern et al. ²⁸	NT	71	62	1.87	0.17	6
Joshi et al. ²⁰	0.48, 0.49, 0.53	39-42	82-86	2.17-3.0	0.68-0.74	9
Blends et al. ⁷	54%	50	47	0.94	1.86	6
Ralphs et al. ²⁹	NT	36	83	2.18	0.76	10



Aorta Palpation

(-)



Study	Reliability	Sensitivity	Specificity	LR+	LR-	QUADAS Score (0-14)
Fink et al. ²⁷	0.66	60	75	2.70	0.43	9
Lederle et al. ¹⁸	NT	50	NT	NT	NT	
Chervu et al. ¹⁵	NT	38	NT	NT	NT	7
		77	NT	NT	NT	
Collis et al. ¹⁶	NT	44	91	5.00	0.62	8
Karkko et al. ²¹	NT	48	NT	NT	NT	4
Kier et al. ²⁴	NT	31	NT	NT	NT	7
Lederle&Sinsel ¹⁷	NT	39	96	12.0	0.72	NA



Appendix Palpation




(-)

Study	Reliability	Sensitivity	Specificity	LR+	LR-	QUADAS Score (0-14)
Campbell & McHaff ¹¹	NT	76	NT	NT	NT	8
Alvarado ¹² (tenderness)	NT	100	13	1.14	0.00	8
Alvarado ¹² (Blumberg's sign)	NT	55	78	2.5	0.58	8
Tzandis et al. ¹³ (tenderness)	NT	90	59	2.19	0.17	9
Tzandis et al. ¹³ (Blumberg's sign)	NT	66	75	2.61	0.45	9
Sola et al. ¹⁴	NT	87	90	8.42	0.15	9


49



Spleen Palpation – Middleton's Maneuver




(-)



Study	Reliability	Sensitivity	Specificity	LR+	LR-	QUADAS Score (0-14)
Tamayo et al. ¹⁷	0.31	35	87	2.69	0.75	11
Barlow et al. ¹⁸	0.7, 0.56, 0.57	56	91	8.00	0.57	9

➤ Are the relationships between the areas on the body chart, the interview, and physical exam consistent? "Do the features fit" a recognizable clinical pattern? **Yes** No

Please explain areas that may need clarification. **R T8/9 CV/CT jt Mechanical Dysfunction**

 Orthopaedic Manual Physical Therapy Series 2017-2018 www.vompti.com


→ Evidence-based Management of Acute Musculoskeletal Pain

Chapter 5 Acute Thoracic Spinal Pain

→ There are currently no guidelines for the management of thoracic spinal pain. This document provides an overview of the evidence in this area to raise awareness of the need for formal population studies on the diagnosis and management of thoracic spinal pain.

Relative Prevalence of Local Causes of Thoracic Pain

Frequency	Entities	Prevalence
Rare conditions	Primary and secondary neoplasia	0.63% (Deyo et al. 1988)
	Disc protrusion	0.15% of all surgically treated disc abnormalities (Love and Schorn 1965)
	Rheumatoid arthritis	Unknown
Uncommon conditions	Spinal infection	< 0.01% (Liang and Komaroff 1982)
	Traumatic fractures	Unknown
Common conditions	Somatic pain	16% of presentations with back pain in primary care (Deyo and Diehl 1988)
	Osteoporotic fractures	6.5% in 50–59 year olds and 77.8% in > 90 year olds (Melton et al. 1989)

 Orthopaedic Manual Physical Therapy Series 2017-2018 www.vompti.com

Thoracic Spine/Ribcage Mechanical Dysfunction – Clinical Characteristics

- Scaringe and Ketner – Manual methods for the treatment of rib dysfunctions and associated functional lesions.
Topics in Clinical Chiropractic (1999)
 - “Costovertebral or costotransverse jt dysfunction will present with localized pain to the posterior thorax that may radiate to the anterior chest or along the associated rib”
 - “Symptoms usually unilateral and painful upon deep inspiration, coughing or sneezing”
 - “Passive or active TL flexion, rotation, and/or lateral flexion may increase the symptoms”
 - “Palpable tenderness of the involved CT jt and rib angle is noted upon joint challenge”
 - “Adjacent thoracic vertebral and rib segments are usually restricted, may complicate the clinical picture, and stimulate or exacerbate protective spasm”



Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test



Thoracic Exam Observation

- General appearance and willingness to move
- Head position
- Posture
 - Cervical curve (presence of lordosis)
 - Thoracic curve (sagittal and frontal planes)
 - Scoliosis (rib hump)
 - Scapular positioning
- Swelling
- Muscle girth and symmetry / changes in body contour
 - Atrophy, spasm, swelling
- Rib movement with breathing
- Skin
 - Scars (especially previous surgical scars)
 - Itchy/redness



Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test



Thoracic Exam



- Screen neighboring joints
 - Cervical motion
 - Rotation with OP
 - Extension
 - Quadrant with OP
 - PAIVM
 - Shoulder functional movement screen
 - Active Elevation, Abduction, Abd/ER, Add/IR with Ops
 - Full Can and ER MMT
 - Passive Quadrant Testing
 - Can be good asterisks and help differentiate between cervical/thoracic/shoulder pathology

(-) adjacent jt clearing



Rib Screening with Respiration (+)

- Deep inhalation and exhalation
 - Rib excursion
 - Upper and lower ribs
 - Quantity
 - Symmetry
 - Pain reproduction
 - May indicate the need to examine the ribs in more detail



Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test






Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Thoracic AROM Assessment

- Thoracic AROM:
 - Flexion
 - Extension
 - Rotation
 - Side bending





Orthopaedic Manual Physical Therapy Series

Combined Motions

(+)


SB → Contra Rotation Rotation → Ipsi SB



 Orthopaedic Manual Physical Therapy Series 2017-2018 www.vompti.com


Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test

 Orthopaedic Manual Physical Therapy Series 2017-2018 www.vompti.com


Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test



Neurodynamics

- Sympathetic chain – anterior along rib heads and CV joint
 - Loaded during flexion, contralateral SB and rotation
 - Further loading with thoracic flexion and contralateral SB in Slump type position (long sitting)
- Critical Zone (T4-9)
 - Narrow, decreased blood supply
 - T6 often considered tension point
 - Segmental stiffness of mid T/S could contribute to signs and symptoms (local and peripheral) associated with adverse neurodynamics
 - Symptoms associated with (+) Slump often are altered after manipulative treatment of mid T/S
 - Possible cord compression – large HNP



Neurodynamic Testing - Sympathetic (Long Sit) Slump Test

- Sympathetic trunk is unilaterally lengthened in the long-sit position (Butler and Slater)
 - More so with contralateral thoracic SB, thoracic rotation, cervical SB
- Often utilized to examine neural tissues in head, neck, thorax and lumbar spine (Butler, 2000)
 - Sympathetic System linked to CRPS II, T4 syndrome, TOS
 - Neurons T1-L2 (head/neck – LE)
 - Recommended when sympathetic trunk is suspected of contributing to symptoms such as hyper or hypohidrosis, altered skin color or temp, or slumped posture mechanism of injury
- Can be position of mobilization and treatment
- Reliability and Validity Unknown
 - Slater et al/Cleland et al – Increased skin conductance and decreased skin temperature following SST



CLINICAL PRESENTATION, QUANTITATIVE SENSORY TESTING, AND THERAPY OF 2 PATIENTS WITH FOURTH THORACIC SYNDROME

Manual Therapy 10 (2005) 292-296

www.

Gary A. Mellick, DO,^a and Larry B. Mellick, MS, MD^b

Case Report

The T4 syndrome

Jenny Louise Conroy, Anthony G. Schneiders*

School of Physiotherapy, University of Otago, P.O. Box 56, Dunedin, New Zealand




Fig 1. Distribution of symptoms of 2 flexion-hyperextension injuries. Case 1 represents an injury from the left side and case 2 represents a posterior impact injury.

Fig. 1. Body chart and symptom behavior. Int = intermittent, sup = superficial.

Practical Applications



- Information from this report will help practitioners better recognize and understand T4 syndrome.
- T4 syndrome should be considered when patients present with a unique constellation of signs and symptoms as described in this report.
- The method of treatment used in this report (injection) may help us better understand the underlying pathomechanics for T4 and other similar syndromes so that similar or other non-invasive therapies may be investigated.



Normal sensory and range of motion (ROM) responses during Thoracic Slump Test (ST) in asymptomatic subjects


Journal of Manual and Manipulative Therapy 2013 VOL. 21 NO. 1

Ketaki C. Joshi, Charu Eapen, Senthil P. Kumar
KMC Mangalore, Manipal University, Mangalore, Karnataka, India




www.vompti.com

Neurodynamic Testing – Slump Test



(-)




Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Neurodynamic Testing - Sympathetic or Long Sit Slump Test

19. SST:
1 and 2:
d lumbar
) Step 3:
al flex-
p 4: add
l lateral
flexion;
sitze by
e flex-
nptoms
duced,
h steps



Orthopaedic Manual Physical Therapy Series 2017-2018 www.vompti.com

Sympathetic (Long Sit) Slump Differentiation



- (+) test defined as reproduction of some or all of the patient's symptoms, asymmetry from uninvolved to involved sides and a (+) sensitizing maneuver
- (+) test suggests sensitivity of the SNS but does NOT indicate that the SNS is the cause of the symptoms or the source of the symptoms


Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test




Practicality??

This Does Exist: PPIVMs/PAIVMs



Thoracic Biomechanical Examination

- Central PA
- Unilateral PA
- Transverse Pressure
 - CT Junction
 - Mid T/S
 - TL Junction
- Upper Thoracic
 - PA like cervical spine
- Mid and Lower Thoracic
 - PA like lumbar spine
- Rib Spring
 - Laterally at rib angles



(+) R T8/9




Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Thoracic – Rib Examination

- Rib Spring
 - Laterally at rib angles



Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Rib Joint Pathomechanics

- Theoretically could be dysfunctional at CV or CT joints
- 1st Rib – often subluxed cranially with trauma or repetitive overuse of UE, as well as TOS
- Commonly have posterior rotation of rib on same side as flexion restriction
 - Thought that restriction of rib movement anteriorly can lead to recurrence of unilateral flexion restriction
- Anterior subluxation
 - Blow to posterior chest wall
 - Prominence of rib anteriorly and concavity posteriorly
- Posterior subluxation
 - Blunt trauma to anterior chest wall



Pathomechanics

- No studies analyzing motion of T/S in subjects with primary or secondary spine disorders (based on anatomy and opinion/models)
- Flexion restrictions (inability of spinal unit to rotate forward in sagittal plane)
 - More common in upper to mid thoracic spine between T3/4 – T6/7 (flattened area and loss of normal posterior kyphosis)
 - Thought to occur after whiplash
 - (+) Flexion, contralateral SB and rotation combined ROM testing
- Extension restrictions (inability of segment to rotate backward in sagittal plane)
 - More common in upper thoracic spine and CT junction C7-T2
 - Also common in lower thoracic spine and TL junction
 - (+) Extension, ipsilateral SB and rotation combined ROM testing



Thoracic Spine/Ribcage Clinical Pearls

- Literature suggests that movement at CV joint creates movement at CT joint and that dysfunctions are rarely specific
- Several authors suggest adjacent thoracic facet, CV and CT joints are often restricted together
- Even though restricted together, treatment directed towards one joint may not result in improvement to other joint
 - Empirical evidence suggests sustained restriction may perpetuate dysfunction if only Facet, CV or CT joint is addressed independently




Thoracic Objective Examination

- Observation/Postural Assessment/Functional Testing
- Cervical shoulder and rib screening
- Thoracic AROM/PROM/Resisted Testing
 - Combined Motions
- Neurological Testing
 - Segmental
 - Central
- Neurodynamic Testing
- Biomechanical Examination
 - Thoracic PAIVM's
 - Rib Spring
- Specific Rib Examination
 - 1st rib CRLF test




Rib Joint Pathomechanics – 1st Rib

- Ant/Middle Scalene insertion
- Lower plexus trunk
- Elevation > Depression
 - Scalene hypertrophy
 - Upper chest breather
 - Prone rotation sleeper
 - Computer/ergonomics
- Potential site of neurovascular compression of plexus, subclavian artery or vein (TOS)
- Hypomobile elevated 1st rib thought to play a role in upper trap symptoms
- “Jump Sign”




Cervical Rotation-Lateral Flexion Test

- Examine mobility of 1st rib
- Pt sitting
- C/S passively and maximally rotated **AWAY** from side being tested
- Gently flex as far as possible, moving ear toward the chest
- (+) if lateral flexion is limited or blocked (+ R and - L)
- Excellent interrater reliability $K = 1.0$ and good agreement with cineradiographic findings $K = .84$



(+)




1st Rib PAIVM Assessment



(+)



Orthopaedic Manual Physical



www.vompti.com

Physical Exam *Asterisks* **Signs/Symptoms** (Special tests, Movement/Joint Dysfunction, Posture, Palpation, etc)

- (-) Cervical and Shoulder Screening
- (+) Thoracic symptoms with deep breath
- Thoracic ROM: (+) Extension, R rotation, R SB, (+) R rot → R SB combined
- (-) Neurological Examination
- (-) Neurodynamic Testing
- (+) T8/9, Rib Spring to R9
- Palpatory changes along angle of R9
- (-) Visceral palpation
- Neck Disability Index = 22% perceived disability



Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

PICO

- In patients with mechanical thoracic spine pain/dysfunction, does the addition of manual therapy help reduce pain and improve function?
- Assessment of current evidence




The screenshot shows the PubMed website interface. At the top is the VOMPTI logo. Below it is the browser address bar showing 'http://www.ncbi.nlm.nih.gov/www.lib...'. The main search area has 'PubMed' selected in the dropdown menu and 'thoracic manipulation' entered in the search box. A dropdown menu shows suggestions: 'thoracic manipulation shoulder', 'thoracic manipulation neck', 'thoracic manipulation neck pain', 'thoracic manipulation and shoulder pain', 'thoracic manipulation shoulder pain', and 'thoracic manipulation cervical'. Below the search area are navigation tabs: 'Using PubMed', 'PubMed Tools', and 'More Resources'. At the bottom of the screenshot, there are two orange bullet points: '• and thoracic pain??' and '• 3 somewhat relevant articles of 1st 100 searched'. The footer of the screenshot includes the VOMPTI logo icon, the text 'Orthopaedic Manual Physical Therapy Series 2017-2018', and the website 'www.vompti.com'.

The Effectiveness of Thoracic Spine Manipulation for the Management of Musculoskeletal Conditions: A Systematic Review and Meta-Analysis of Randomized Clinical Trials

RONALD F. WALSER, PT, DPT¹; BRENT B. MESERVE, PT, DPT²; THOMAS R. BOUCHER, PhD³

- 13 studies analyzed (RCTs) – 3 for shoulder, 9 for cervical conditions, 1 on lower trap function
- Identified need for additional studies to examine effectiveness of TSM
- NO studies investigated effect of thoracic spine manipulation on thoracic spine symptoms


Orthopaedic Manual Physical Therapy Series 2017-2018
www.vompti.com

→ Evidence-based Management of Acute Musculoskeletal Pain

Chapter

5

Acute Thoracic Spinal Pain

→ There are currently no guidelines for the management of thoracic spinal pain.

Evidence of Benefit


Manual Treatment

There have been no systematic reviews of therapy for thoracic spinal pain. Schiller (2001) compared the use of spinal manipulation with non-functional ultrasound placebo in a small, randomised controlled trial of 30 patients with mechanical thoracic spinal pain. This demonstrated significantly better reductions in numerical pain ratings and improvements in lateral flexion with manipulation at the end of a two to three week treatment period. These changes were maintained a month later, but were no longer better than in the placebo group. Notably there were no significant differences in McGill pain questionnaires and Oswestry Back Disability Indices between groups at any point in the trial. The small sample size was suggested as a reason for this, leaving unanswered questions about the real efficacy of manipulation.

Other Treatment

No studies can be found that address the treatment of acute thoracic spinal pain with the following therapies:

- consumer education
- reassurance and home rehabilitation
- drug therapy
- bed rest
- mobilisation
- functional restoration
- behavioural therapy
- back school
- exercises
- injection treatments
- surgery

INTERVENTIONS	EVIDENCE LEVEL
<div style="display: flex; align-items: center; margin-bottom: 5px;">  <div style="margin-left: 5px;"> <p style="font-size: small; margin: 0;">Evidence of Benefit</p> <p style="font-size: x-small; margin: 0;"><i>Spinal Manipulation</i> — There is evidence from one small study that spinal manipulation is effective compared to placebo in thoracic spinal pain.</p> </div> </div>	<p style="font-size: x-small; margin: 0;">LEVEL II: Schiller 2001</p>

com

Differential Diagnosis and Treatment in a Patient With Posterior Upper Thoracic Pain

Stacie J Fruth
PHYS THER. 2006; 86:254-268.

- Case study 35 y/o male with 4 month hx of symptoms
- Multifactorial manual therapy approach
- Discusses CV/CT joint assessment – 2 separate joints but assessed together due to proximity and shared movement with function
- Differential Diagnosis ruling out other musculoskeletal and visceral sources

Rationale for Treatment

According to Scaringe and Ketner⁴ and Triano et al,¹⁰ treatment of CV and CT joint dysfunction should include attempts to normalize mechanics by soft tissue and joint mobilization or manipulation, scapular stabilization and postural reeducation, and any necessary pain control measures. Based on this recommendation and


zation and postural exercises. I was unable to find any studies that examined the effects of joint mobilization on either the thoracic spine or the CV and CT joints.



Orthopaedic Man

www.vompti.com

Thoracic Treatment - Mobilization




Central PA

Unilateral PA

Transverse Pressures



Rotary PA Facet




Rotary PA Rib



Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Prone Rotary PA HVLAT (Facet T2-9 vs. R2-9 Costotransverse)




Costotransverse



Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Thoracic Spine Extension Mobilization with Foam Roller



Seated Mid Thoracic Distraction




Figure 1. Seated thoracic spine distraction thrust manipulation used in this study. The therapist uses his or her sternum as a fulcrum on the subject's middle thoracic spine and applies a high-velocity distraction thrust in an upward direction.



oti.com

Supine Upper and Mid-Thoracic AP HVLAT



m

Alternate Thoracic and Ribcage Techniques


- Flexion Bias (T3/4 – T6/7)
- Extension Bias (CT Junc and TL Junc)
- Rib Manipulation
 - Exhalation (SB towards lesion, ½ breath in and breath out)




Mobilization in Sympathetic (Long-Sit) Slump Position



ti.com



1st Rib Manipulation: “Snooker” Technique



Orthopaedic Manual Physical Therapy Series

Common OCS Question - Rib Subluxation MET



Figure 22. Posterior subluxed right sixth rib treatment using a muscle energy technique. Reprinted from Orthopaedic Manual Physical Therapy Management of the Cervical-Thoracic Spine and Ribcage. Copyright 2000, with permission of the publisher Evidence in Motion, LLC (www.evidenceinmotion.com).



Figure 21. Anterior subluxed right sixth rib treatment using a muscle energy procedure. Reprinted from Orthopaedic Manual Physical Therapy Management of the Cervical-Thoracic Spine and Ribcage. Copyright 2000, with permission of the publisher Evidence in Motion, LLC (www.evidenceinmotion.com).



Orthopaedic Manual Physical Therapy Series 2017-2018

www.vompti.com

Pattern Recognition

Identify the key subjective and physical features (i.e. **clinical pattern**) that would help you recognize this disorder in the future.

Subjective	Physical
<p>Mechanical nature of symptoms Somatic quality, even of referral Sensitivity to deep breath, twist, cough/sneeze (ribcage) Ergonomic/work posture component</p>	<p>Pain upon T/S ROM testing (Rot, SB, Ext) Palpatory changes along rib angle (-) Neurodynamic testing Associated dysfunction of same segment Z Jt Restriction and/or symptoms with PAIVM/PPIVM testing</p>

